

Mirage Cosmology on Unstable D3-Brane Universe

Anastasios Psinas¹

Department of Physics, Northeastern University, Boston, MA 02115-5000, USA

E-mail: psinas.a@neu.edu

Abstract. We study the geodesic motion of an unstable brane moving in a higher dimensional bulk spacetime. The tachyon which is coupled to a $U(1)$ gauge field induces a non-trivial cosmological evolution. Interestingly enough, this system exhibits a much smoother initial cosmological singularity in comparison with former works.

String cosmology has been one of the leading approaches towards describing cosmos at large scale. Based on the very rich structure of string theory and its dualities cosmologists have acquired a better understanding of the evolution of the universe. In particular, the progress made in the AdS/CFT [1] correspondence gave significant impetus towards the development of various cosmological models. In one of those [2], the universe is described by a Dp -brane (with $p = 3$ for a 4-dim universe) moving in a higher p' dimensional space. According to this, the ambient space is filled by a stack of Dp' -branes with $p' > p$ which is much heavier than the probe so that the back-reaction is neglected. The cosmological expansion or contraction that these universes exhibit can be ascribed to mirage matter that may come from the bulk. Such constructions have very interesting consequences with respect to the problem of the initial singularity. For instance, one can study the effects of different kind of matter on the effective energy density of the probe brane by investigating by how much and to what extent cosmological singularities are regularized. Essentially, the energy content of the theory can give us details of the spacetime structure on the brane universe.

The instabilities in the spectrum of string theory (in the form of tachyons) have been a serious issue for quite some time. Quite recently though, it is shown, that unstable D-branes can form and then decay so that we end up with a stable string vacuum [3, 4]. It is therefore believed that the very early history of the universe was dominated by instabilities of tachyonic nature. Several implementations of the tachyon in cosmology are known. In [4] there has been an extensive analysis of mirage cosmology for tachyons which live in the bulk. Models of unstable universes are also obtained in the case of the geodesic motion of a tachyonic Dp -brane for which the tachyon field is confined on the defect [6]. Thinking in the same spirit, it would be instructive to examine the possibility of having tachyon matter coupled to an abelian gauge field where both of them reside on the brane [7]. Keep in mind that unlike the brane-antibrane models [8] we just consider one mobile D3-brane while the gauge group is $U(1)$. We also expect the gauge degrees of freedom to play an important role at early times. For instance, we recall that even in the FRW model the state equation of the cosmic fluid is that of a photonic gas during the

¹ Present address: Department of Physics, University of Patras, GR-26500, Patras, GREECE. Talk presented at the RTN meeting “The Quest for Unification: Theory Confronts Experiment”, Corfu, Greece, Sept, 11-18, 2005

first stages of the evolution of cosmos. In fact, as it will be shown, our analysis clearly points out that at very early times the universe is radiation dominated.

We start with the most general ansatz of a ten dimensional static and spherically symmetric spacetime

$$ds_{10}^2 = g_{00}(r)dt^2 + g(r)(d\vec{x})^2 + g_{rr}dr^2 + g_S(r)d\Omega_5 \quad (1)$$

where the timelike part of the metric g_{00} is negative. Additionally, the complex tachyon on the D3-brane couples to the massless gauge field in the following fashion

$$S_3 = -T_3 \int d^4\xi e^{-\phi} V(T) \sqrt{-\det \tilde{K}_{\mu\nu}} - T_3 \int d^4\xi \hat{C}_4 \quad (2)$$

where

$$\tilde{K}_{\mu\nu} = K_{MN} \frac{\partial x^M}{\partial x^\mu} \frac{\partial x^N}{\partial x^\nu} \quad (3)$$

and

$$K_{MN} = g_{MN} + 2\pi\alpha' F_{MN} + \frac{1}{2}(D_M T D_N T^* + c.c) \quad (4)$$

We ascribe capital indices M,N for the bulk coordinates while μ, ν are the one kept for the probe brane. We note that $F_{\mu\nu}$ is the world volume antisymmetric gauge field strength and g_{MN} is the induced metric on the D3-brane universe and ϕ is the dilaton. Observe, that the RR field $\hat{C}(r) = C_{0\dots 3}(r)$ is also included in the action. In addition, the form of the complex tachyon field and its covariant derivative are given as follows

$$D_M T = \partial_M T - i A_M T \quad (5)$$

$$T = T_x + iT_y \quad (6)$$

Here we underline that for the rest of the analysis we will consider that $dV(T)/dT \simeq 0$. This is equivalent to say that we are in the regime of small T . In other words we are mainly focused on exploring the dynamics of the D-brane at a point which the instability has started to occur.

In our analysis we assign an expectation value to the tachyon, i.e. $T_y = c$ while we adopt the gauge $A_0 = 0$. The remaining non-vanishing spatial components A_i define the following combination

$$\alpha^i = A_0^i c \quad (7)$$

It is crucial to mention that we study small and time-dependent fluctuations of the gauge field around A_0^i which is taken to be small. All of the above translate to

$$A^i(t) = A_0^i + \tilde{A}^i(t) \quad (8)$$

Thus our Lagrangian after some obvious redefinitions turns out to be

$$L = -\sqrt{M - B\dot{r}^2 - Ch_{ij}\dot{\varphi}^i\dot{\varphi}^j - D\dot{T}^2 - E(\alpha^i\dot{A}_i\dot{T} + \dot{A}_i\dot{A}^i)} + \hat{C}(r) \quad (9)$$

The presence of the radical in the lagrangian will certainly complicate the form of the equation of motions of the associated fields [7]. Expanding the square root is a good approximation only at late times since for small scale factors $a(r(t))$ the above approximation scheme fails to capture the underlying physics at the $a \rightarrow 0$ limit. Besides, the tachyon induced instabilities will be more prominent as $a \rightarrow 0$ since at very late times the tachyon will condense. In addition, it would be interesting to examine the above system for small scale factors so that we improve our understanding off the tachyon-gauge field-gravity interactions at high curvature.

Interestingly enough, the induced metric on the brane describes a flat FRW spacetime which reads

$$d\hat{s}^2 = -d\eta^2 + g(r(\eta))(d\vec{x})^2 \quad (10)$$

while the cosmic time is defined as

$$d\eta = \frac{|g_{00}|}{\mathcal{E}} \sqrt{T_3^2 V_0^2 e^{-2\phi} g^3 + Q_i Q_i + \frac{(2g\tilde{Q} - \alpha_i Q_i)^2}{(4g^2 - \alpha_i \alpha_i)}} dt \quad (11)$$

For the sake of clarity we note that $\mathcal{E}$ denotes the energy of the system while Q_i and $\tilde{Q}$ stand for the integration constants of the gauge field and the tachyon respectively. Finally, due to the spherical symmetry of the ambient space the angular momentum l is conserved.

Close inspection of Eq. (11) reveals that the argument of the radical is not only non-positive but it is also singular as well when $4g^2(r) = \alpha_i \alpha_i$ unlike in models like [2]. Nonetheless, we are primarily interested in the regime of very small scale factors. At that limit, it is easy to check, that the argument in the radical is strictly nonnegative (for nonzero values of the Q_i charges) while the singularity has been avoided. Our analysis so far pertains a very general class of metrics (equipped with certain symmetries) describing the bulk. A very characteristic example of a spacetime of this sort, is the well studied case of the near-horizon limit of a black hole in an $AdS_5 \times S^5$ background

$$ds^2 = \frac{r^2}{L^2} (-f(r)dt^2 + (d\vec{x})^2) + \frac{L^2}{r^2} \frac{dr^2}{f(r)} + L^2 d\Omega_5^2 \quad (12)$$

where $f(r) = 1 - \frac{r_0^4}{r^4}$ and $\hat{C}(r) = \frac{r^4}{L^4} - \frac{r_0^4}{2L^4}$

Given the very simple form of the induced metric on the brane, one can straightforwardly derive the FRW equations. Thus, the effective energy density on the probe reads as follows

$$\frac{8\pi G}{3} \rho_{\text{eff}} = \frac{1}{a^2 L^2 F^2} ((\mathcal{E} + a^4)^2 - a^2 (1 - \frac{r_0^4}{L^4 a^4}) (F^2 + \frac{l_S^2}{L^2})) \quad (13)$$

The constant term accompanying the RR term is absorbed in $\mathcal{E}$. Also, F is given as a function of the scale factor of the universe in the following form

$$F^2 = T_3^2 V_0^2 e^{-2\phi} a^6 + Q_i Q_i + \frac{(2\tilde{Q}a^2 - \alpha_i Q_i)^2}{4a^4 - \alpha_i \alpha_i} \quad (14)$$

In principle, the singularities a given spacetime exhibits can be related to the energy content of the theory living in a given background. Thus, it is reasonable to expect that the blowing up of the energy density on the brane is indicative of a curvature (initial) singularity. Thus, we are interested in exploring the behavior of ρ_{eff} as $a \rightarrow 0$.²

Let us now study in detail the behavior of the energy density in the limit of very small scale factors. It turns out, that as $a \rightarrow 0$ F^2 reads as follows

$$F^2 = T_3^2 V_0^2 e^{-2\phi} a^6 + |\vec{Q}|^2 \sin^2 \theta \quad (15)$$

where θ denotes the angle between vectors α_i and Q_i . One observes that ρ_{eff} has no explicit dependance on α_i which manifests the non trivial interaction between the tachyon and the gauge field. However, the interaction appears implicitly through the $|\vec{Q}|^2 \sin^2 \theta$ term. Also, note that whenever $\sin \theta$ becomes zero the effective energy density on the unstable probe may be similar

² An alternative method of characterizing a singularity as a curvature singularity is through the evaluation of the Kretschmann scalar $R_{\mu\nu\kappa\lambda} R^{\mu\nu\kappa\lambda}$. However in the context of our analysis this method is not preferable.

to the one of a stable D3-brane[2]. In all other cases this novel effect doesn't occur due to the non vanishing integration constants. We also stress, that the singular behavior that the F^2 term exhibited in the most general case, has disappeared at the very early stages of the evolution of the universe.

A simpler version of Eq. (13) is obtained when the gauge field vanishes. It turns out that in this case the presence of tachyon regulates the cosmological evolution by smoothening the initial singularity. In essence, our analysis corroborates former results found in [6] for a vanishing Ramond-Ramond field.

For the sake of concreteness we denote the full expressions of the energy density and effective pressure on the brane universe in the $a \rightarrow 0$ limit. To this end, we express ρ_{eff} and p_{eff} in powers of inverse scale factor so that it is easier to read the value of w which enters in the state equation $p_{\text{eff}} = w\rho_{\text{eff}}$ of the cosmic fluid. Thus, we obtain

$$\begin{aligned}
p_{\text{eff}} = & \frac{1}{8\pi GL^2} \frac{r_0^4}{L^4} \left(1 + \frac{l^2}{L^2} \frac{1}{Q_i^2 \sin^2 \theta}\right) \frac{1}{a^4} \\
& + \left(\frac{\mathcal{E}^2}{8\pi GL^2}\right) \left(-\frac{1}{(Q_i^2 \sin^2 \theta)^2} + \frac{8r_0^4}{L^4} \frac{l^2}{L^2} \frac{\alpha_i Q_i \tilde{Q}}{\alpha_i^2 (Q_i^2 \sin^2 \theta)^2}\right) \frac{1}{a^2} \\
& + \frac{3}{8\pi GL^2} \left(1 + \frac{l^2}{L^2} \frac{1}{Q_i^2 \sin^2 \theta} + \frac{8\mathcal{E}^2}{3} \frac{\alpha_i Q_i \tilde{Q}}{\alpha_i^2 (Q_i^2 \sin^2 \theta)^2} - \frac{16}{3} \frac{r_0^4}{L^4} \frac{l^2}{L^2} \frac{(\alpha_i Q_i)^2}{\alpha_i^4}\right)
\end{aligned} \tag{16}$$

while the effective density is given by

$$\rho_{\text{eff}} = \frac{3}{8\pi GL^2} \frac{r_0^4}{L^4} \left(1 + \frac{l^2}{L^2} \frac{1}{Q_i^2 \sin^2 \theta}\right) \frac{1}{a^4} + \frac{3\mathcal{E}^2}{8\pi GL^2} \frac{1}{(Q_i^2 \sin^2 \theta)^2} \frac{1}{a^2} - \frac{3}{8\pi GL^2} \left(1 + \frac{l^2}{L^2} \frac{1}{Q_i^2 \sin^2 \theta}\right) \tag{17}$$

At this point, let us comment on Eq. (16,17) in detail. First of all, it is clear that the energy density has no explicit dependance on the tachyon integration constant $\tilde{Q}$ while only the gauge field constants Q_i survive. It looks as if the tachyonic degrees of freedom have no effect on ρ_{eff} as $a \rightarrow 0$ when a gauge field-tachyon interaction is present. However, this is not the case with the effective pressure on the brane. In addition, it turns out that $w = 1/3$ at the very early history of the universe leading one to the conclusion that the universe is radiation dominated. Thus, initially the gauge field prevails over the tachyon. At later evolutionary stages w depends on the values of the parameters of the system (Q_i , $\tilde{Q}$, α_i , $\sin \theta$, $\mathcal{E}$, l , r_0 , L) and so causality may be violated. Closer inspection of Eq. (16,17) reveals that at late times a fine tuning of the parameters can lead to a pressureless universe with a finite energy density which could be negative. The negativity of the energy density is a manifestation of the tachyon.

An equally interesting case is the one with a vanishing α_i . One can easily show that w takes the following values $w = [\frac{1}{3}, -\frac{1}{3}, -1]$. Initially, the probe is radiation dominated however at later times it has the state of a universe filled with a cosmic fluid that resembles a non-vanishing cosmological constant. So, even though $w = -1$ the associated energy density is negative due to the tachyon. Interestingly enough, in this case, $|w| \leq 1$ and thus causality is respected. The above values for w are reminiscent of quintessence in which the accelerated expansion of the universe occurs when $-1 < w < -\frac{1}{3}$ [9]. For completeness, the term $\sim a^{-2}$ in ρ_{eff} has the same effect as a negative curvature term responsible for accelerating the expansion of the universe.

The effects of the minimal tachyon-gauged boson coupling on the initial cosmological singularity are quite transparent. According to Eq. (17) the energy density is not that divergent as $a \rightarrow 0$ in comparison to former works [2]. More specifically, in the model at hand, the most divergent term of the energy density goes like $\sim a^{-4}$ while for stable D-branes [2] it is shown

that $\rho_{\text{eff}} \sim a^{-8}$. This in particular leads one to the conclusion that the above coupling makes the singularity of spacetime less severe. For purely tachyonic branes a similar result was initially reported in [6].

In our approach we were mainly interested in dealing with unstable D3-branes moving in a geodesic embedded in a higher dimensional bulk. Although in models like ours it is very hard to obtain solutions in closed form we didn't face such a problem. This can be attributed to the fact that we were focused in a region where the tachyon potential was very flat. The presence of the minimal gauge field-tachyon coupling was shown to have significant cosmological consequences. According to our analysis, the universe is proven to be radiation dominated at its incipient stages while at later epochs it can evolve in a more complicated fashion. This behavior clearly shows that the gauge degrees of freedom are prevalent, however as the universe evolves the effects of the tachyon become more apparent. Further, it is shown, that the initial singularity which the mirage models exhibit is softened quite significantly. This particular result looks quite promising even though we were not able to completely remove the singularity. We hope that further work in this direction will provide more clues on how to tame cosmological singularities.

Acknowledgements

I would like to thank the organizers of the RTN meetings "The Quest for Unification: Theory Confronts Experiment" and "Constituents, Fundamental Forces, and Symmetries of the Universe" held in Corfu, Greece, September 11-26, 2005, for financial support and for providing me the opportunity to present the results of my work.

- [1] Maldacena J M 1997 The large N limit of superconformal field theories and supergravity *Adv. Theor. Math. Phys.* **2**, 231 (1998), *Int. J. Theor. Phys.* **38**, 1113 (1999) (*Preprint* arXiv:hep-th/9711200)
- [2] Kehagias A, Kiritsis E 1999 Mirage Cosmology *JHEP* **9911**, 022 (1999) (*Preprint* hep-th/9910174)
- [3] Garousi M R 2000 Tachyon couplings on non-BPS D-branes and Dirac-Born-Infeld action *Nucl. Phys. B* **584**, 284 (2000) (*Preprint* hep-th/0003122)
- Bergshoeff E A, de Roo M, de Wit T C, Eyras E Panda S 2000 T-duality and actions for non-BPS D-branes *JHEP* **0005**, 009 (2000) (*Preprint* hep-th/0003221)
- Mazumbar A, Panda S, Perez-Lorenzana A 2001 Assisted inflation via tachyon condensation *Nucl. Phys. B* **614**, 101 (2001) (*Preprint* hep-ph/0107058)
- Kluson J 2000 Proposal for non-BPS D-brane action *Phys. Rev. D* **62**, 126003 (2000) (*Preprint* hep-th/0004106)
- Choudhury D, Ghoshal D, Jatkar D P, Panda S 2002 On the cosmological relevance of the tachyon *Phys. Lett. B* **544**, 231 (2002) (*Preprint* hep-th/0204204)
- [4] Sen A 2002 Rolling tachyon *JHEP* **0204**, 048 (2002) (*Preprint* hep-th/0203211)
- Sen A 2002 Tachyon matter *JHEP* **0207**, 065 (2002) (*Preprint* hep-th/0203265)
- Sen A 2002 Field theory of tachyon matter *Mod. Phys. Lett. A* **17**, 1797 (2002) (*Preprint* hep-th/0204143)
- [5] Kim J Y 2001 Brane inflation in tachyonic and non-tachyonic type 0B string theories *Phys. Rev. D* **63**, 045014 (2001) (*Preprint* hep-th/0009175)
- Papantonopoulos E 2001 Cosmological evolution of a brane universe in a type 0 string background *Phys. Rev. D* **63**, 103506 (2001) (*Preprint* hep-th/0010014)
- Diamandis G A, Georgalas B C, Mavromatos N E, Papantonopoulos E 2002 Acceleration of the universe in type-0 non-critical strings *Int. J. Mod. Phys. A* **17**, 4567 (2002) (*Preprint* hep-th/0203241)
- [6] Jeong D H, Kim J Y 2005 Mirage cosmology with an unstable probe D3-brane *Phys. Rev. D* **72**, 087503 (2005) (*Preprint* hep-th/0502130)
- [7] Psinas A 2006 Mirage cosmology of U(1) gauge field on unstable D3 brane universe *JHEP* **0601**, 138 (2006) (*Preprint* hep-th/0509001)
- [8] Barnaby N Cline J M 2004 Creating the universe from brane-antibrane annihilation *Phys. Rev. D* **70**, 023506 (2004) (*Preprint* hep-th/0403223)
- Sen A 2003 Dirac-Born-Infeld action on the tachyon kink and vortex *Phys. Rev. D* **68**, 066008 (2003) (*Preprint* hep-th/0303057)
- [9] Hellerman S, Kaloper N, Susskind L 2001 String theory and quintessence *JHEP* **0106**, 003 (2001) (*Preprint* hep-th/0104180)

Fischler W, Kashani-Poor A, McNees R, Paban S 2001 The acceleration of the universe, a challenge for string theory *JHEP* **0107**, 003 (2001) (*Preprint* hep-th/0104181)